



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$=r\sin\theta+\sqrt{[r^2\sin^2\theta-2rx(1+\cos\theta)]}.$$

$$\therefore x=\frac{2r[(\sin\theta-\sin\varphi)\sin(\theta+\varphi)-(\sin\theta-\sin\varphi)^2]}{(\cos\theta+\cos\varphi)^2}$$

$$=2r[\sin\frac{1}{2}(\theta+\varphi)\sin\frac{1}{2}(\theta-\varphi)\sec^2\frac{1}{2}(\theta-\varphi)-\tan^2\frac{1}{2}(\theta-\varphi)].$$

$$\pi x^2=4\pi r^2[\sin\frac{1}{2}(\theta+\varphi)\sin\frac{1}{2}(\theta-\varphi)\sec^2\frac{1}{2}(\theta-\varphi)-\tan^2\frac{1}{2}(\theta-\varphi)]^2.$$

Let L =average length, Δ =average area.

$$\therefore L=\frac{\int_0^\pi \int_0^\theta x d\theta d\varphi}{\int_0^\pi \int_0^\theta d\theta d\varphi}=\frac{2}{\pi^2}\int_0^\pi \int_0^\theta x d\theta d\varphi$$

$$=\frac{8r}{\pi^2}\int_0^\pi(\theta\cos^2\frac{1}{2}\theta-\sin\theta+\sin\theta\log\sec\frac{1}{2}\theta)d\theta=\frac{2r}{\pi^2}(\pi^2-8)=.3789r.$$

$$\Delta=\frac{\pi \int_0^\pi \int_0^\theta x^2 d\theta d\varphi}{\int_0^\pi \int_0^\theta d\theta d\varphi}=\frac{2}{\pi}\int_0^\pi \int_0^\theta x^2 d\theta d\varphi$$

$$=\frac{32r^2}{3\pi}\int_0^\pi(6\theta\cos^4\frac{1}{2}\theta-3\theta\cos^2\frac{1}{2}\theta-6\sin\frac{1}{2}\theta\cos^3\frac{1}{2}\theta+2\sin^3\frac{1}{2}\theta\cos\frac{1}{2}\theta$$

$$-12\sin\frac{1}{2}\theta\cos^3\frac{1}{2}\theta\log\cos\frac{1}{2}\theta)d\theta=\frac{4r^2}{3\pi}(3\pi^2-28)=.2174\pi r^2.$$

85. Proposed by G. B. M. ZERR, A. M., Ph. D., Professor of Science and Mathematics, Chester High School, Chester, Pa.

Two points are taken at random in a circle and a chord drawn through them; a point is then taken at random in each segment. Find the average area of the quadrilateral formed by joining the four points.

Solution by the PROPOSER.

Let P, Q, R, S be the four random points taken as indicated in the problem, NN' the chord through PQ , MM', TT' the chords through S, R , respectively.

Draw CD perpendicular and CD' parallel to NN' . Let $CD=r$, $CE=u$, $CF=v$, $CG=w$, $NQ=x$, $PQ=y$, $NF=\sqrt{(r^2-v^2)}=z$, $TG=\sqrt{(r^2-w^2)}=t$, $ME=\sqrt{(r^2-u^2)}=s$, $\angle D'CA'=\theta$.

An element of the circle at P is $dvdx$; at Q , $y d\theta dy$; at R , $2tdw$; at S , $2sdu$.

